MEDICATION DISCREPANCY TOOL (MDT)

MDT is designed to facilitate reconciliation of medication regimen across settings and prescribers

Medication Discrepancy Event Description: Complete one form for each discrepancy Causes and Contributing Factors :: Check all that apply :: Italicized text suggests patient's perspective and/or intended meaning Patient Level -1.
Adverse Drug Reaction or side effects 6. Intentional non-adherence "I was told to take this but I choose not to." 2. \Box Intolerance 7. Non-intentional non-adherence (ie: Knowledge deficit) 3. Didn't fill prescription "I don't understand how to take this medication." 4. Didn't need prescription 8. Performance deficit 5.
Money/financial barriers "Maybe someone showed me, but I can't demonstrate to you that I can." System Level 13. \Box Duplication. Taking multiple drugs with the same action 10.
Conflicting information from different without any rationale. informational sources 14. \Box Incorrect dosage For example, discharge instructions indicate one thing and pill bottle says another. 15. \Box Incorrect quantity 11. \Box Confusion between brand & generic names 16. \Box Incorrect label 12. Discharge instructions 17. \Box Cognitive impairment not recognized incomplete/inaccurate/illegible 18. \Box No caregiver/need for assistance not recognized Either the patient cannot make out the 19.
Sight/dexterity limitations not recognized hand- writing or the information is not written in lay terms. \checkmark Resolution :: check all that apply Discussed potential benefits and harm that may result from non-adherence □ Encouraged patient to call PCP/specialist about problem Encouraged patient to schedule an appointment with PCP/specialist to discuss problem at next visit Encouraged patient to talk to pharmacist about problem □ Addressed performance/knowledge deficit □ Provided resource information to facilitate adherence Other